

Clinical features

Any acute onset of hypotension or bronchospasm or upper airway obstruction where anaphylaxis is considered possible, **even if typical skin features are not present**

OR


Any acute onset illness with typical skin features (urticarial rash or erythema/flushing, and/or angioedema)

PLUS

Involvement of respiratory, cardiovascular, or persistent severe gastrointestinal symptoms

1 Immediate action

- Call for assistance.
- Lay the patient flat – do not allow them to stand or walk. If unconscious place in recovery position and maintain airway. If breathing is difficult allow the patient to sit (as illustrated).



2 Give INTRAMUSCULAR ADRENALINE (EPINEPHRINE) into mid-lateral thigh without delay

Adrenaline dose chart (1:1000 ampoules containing 1 mg adrenaline per 1 mL)

Age (years)	Weight (kg)	Adrenaline volume 1:1000
<1	5–10	0.05–0.1 mL
1–2	10	0.1 mL
2–3	15	0.15 mL
4–6	20	0.2 mL
7–10	30	0.3 mL
10–12	40	0.4 mL
>12 and adult	>50	0.5 mL

Autoinjector

An adrenaline autoinjector, e.g. EpiPen, may be used instead of an adrenaline ampoule and syringe. For children 10–20 kg (aged 1–5 years) a 0.15 mg device, e.g. EpiPen Jr, should be used. Instructions are on device labels.

Repeat adrenaline every 5 minutes as needed

If multiple doses are required, consider adrenaline infusion if skills and equipment available (see step 5).

Remove allergen (if still present): flick out insect stings, freeze ticks with liquid nitrogen or ether-containing spray (if available) and allow to drop off.

ALWAYS give adrenaline FIRST, then asthma reliever puffer, if someone with known asthma and allergy to food, insects or medicine has SUDDEN BREATHING DIFFICULTY (including wheeze, persistent cough or hoarse voice) even if there are no skin symptoms.

3 Call ambulance to transport patient to hospital

Keep the patient flat and transfer to ambulance via stretcher. Do not allow them to stand or walk even if they appear to have recovered following administration of adrenaline.

4 Supportive management

When skills and equipment are available:

- monitor pulse, blood pressure, respiratory rate, pulse oximetry
- give oxygen and airway support if needed
- obtain intravenous access in adults and hypotensive children
- if hypotensive, give intravenous normal saline (20 mL/kg rapidly) and consider additional wide-bore intravenous access.

5 Additional measures

Adrenaline (epinephrine) infusion

If inadequate response or deterioration, start an intravenous adrenaline infusion as follows:

Give only in liaison with an appropriate specialist. Phone

- Mix 1 mL of 1:1000 adrenaline in 1000 mL of normal saline
- Start infusion at 5 mL/kg/hour (0.1 microgram/kg/min)
- Titrate rate according to response
- Monitor continuously

If adrenaline (epinephrine) infusion is ineffective or unavailable, also consider:

For upper airway obstruction

- nebulised adrenaline (5 mL, i.e. 5 ampoules of 1:1000)
- intubation if skills and equipment are available

For persistent hypotension/shock

- give normal saline (maximum 50 mL/kg in the first 30 min)
- in patients with cardiogenic shock (especially if taking beta blockers) consider an intravenous glucagon bolus of 1–2 mg in adults (in children: 20–30 micrograms/kg up to 1 mg). This may be repeated or followed by an infusion of 1–2 mg/hour in adults
- in adults, selective vasoconstrictors metaraminol (2–10 mg) or argipressin (vasopressin) (10–40 units) only after advice from an appropriate specialist

For persistent wheeze

- bronchodilators: salbutamol 8–12 puffs of 100 micrograms using a spacer or 5 mg salbutamol by nebuliser
- oral prednisolone 1 mg/kg (maximum 50 mg) or intravenous hydrocortisone 5 mg/kg (maximum 200 mg)

6 Observation

Prolonged and biphasic reactions may occur.

Observe the patient for at least 4 hours after last dose of adrenaline.

Observe longer (overnight) if the patient:

- had a severe reaction (hypotension or hypoxia), or
- required repeated doses of adrenaline, or
- has a history of asthma or protracted anaphylaxis, or
- has other concomitant illness, or
- lives alone or is remote from medical care, or
- has known systemic mastocytosis.

Document food, medicine, sting/bite exposure in the 2–4 hours before anaphylaxis.

7 Follow-up treatment

Corticosteroids

The role of corticosteroids is unknown. It is reasonable to prescribe a 2-day course of oral steroid (e.g. prednisolone 1 mg/kg, maximum 50 mg daily) to reduce the risk of symptom recurrence after a severe reaction or a reaction with marked or persistent wheeze. Corticosteroids should only be administered after adrenaline and resuscitation.

Adrenaline (epinephrine) autoinjector

Prescribe an autoinjector, pending specialist review. Train the patient in autoinjector use and give them an ASCIA Action Plan for Anaphylaxis - www.allergy.org.au.

Allergy specialist

Refer patients with anaphylaxis for review.

Antihistamines

Antihistamines have no role in treating respiratory or cardiovascular symptoms of anaphylaxis. Oral non-sedating antihistamines treat itch and urticaria. Injectable promethazine should NOT be used in anaphylactic shock as it can worsen hypotension.