Table: Means and standard deviation of the SIAB-EX (worst) for English speaking eating disordered probands

Item No.		Total group N=351 M (SD)	Anorexia Nervosa restrictive type N=102 M (SD)	Anorexia Nervosa binging/purging type N=121 M (SD)	Bulimia Nervosa N=38 M (SD)	Eating Disorder Not Otherwise Specified N=90 M (SD)
	I Body Image and Slimness Ideal (BI)	**2.6 (0.6)**	**2.7 (0.5)**	**2.9 (0.4)**	**2.0 (0.4)**	**2.2 (0.6)**
16	Body image disturbance	3.2 (1.0)	3.4 (0.8)	3.5 (0.7)	2.3 (1.0)	2.8 (1.0)
7	Fear of gaining weight or getting fat – weight phobia	3.1 (0.8)	3.3 (0.7)	3.5 (0.7)	2.7 (0.8)	2.8 (0.8)
61	Preoccupation with body slimness, figure and body weight	3.5 (0.8)	3.5 (0.6)	3.7 (0.6)	3.4 (0.8)	3.0 (1.0)
(1)	Underweight (modified) / deviance from normal body weight	2.6 (1.3)	3.3 (0.9)	3.2 (0.9)	0.6 (0.7)	2.0 (1.2)
12	Qualitative food reduction	3.1 (1.1)	3.2 (1.1)	3.4 (1.0)	2.7 (1.0)	2.9 (1.1)
59	Compulsive behavior concerning food or eating	3.0 (1.0)	3.1 (0.9)	3.4 (0.8)	2.8 (1.0)	2.5 (1.0)
42	Excessive exercise	2.9 (1.2)	2.8 (1.3)	3.0 (1.2)	2.7 (1.1)	2.8 (1.1)
60	Preoccupation with food or eating	3.5 (0.8)	3.6 (0.7)	3.8 (0.5)	3.5 (0.8)	3.0 (1.0)
14	Chewing and spitting out food	0.9 (1.4)	0.8 (1.4)	1.4 (1.6)	0.9 (1.1)	0.5 (1.1)
3	Internal achivement orientation	2.5 (1.1)	2.7 (1.1)	2.7 (1.0)	1.9 (1.1)	2.2 (1.1)
10	Dependence of self-esteem on figure and weight	3.4 (0.8)	3.4 (0.8)	3.5 (0.7)	3.2 (0.8)	3.2 (0.7)
13	Limits for caloric intake	3.6 (0.9)	3.5 (0.9)	3.8 (0.5)	3.3 (1.1)	3.5 (1.0)
19	Amenorrhea (modified)	1.5 (0.9)	1.9 (0.7)	2.0 (0.5)	1.0 (0.9)	0.6 (0.8)
55	Constipation	1.9 (1.7)	1.8 (1.6)	2.3 (1.7)	1.2 (1.4)	1.6 (1.6)
15	Regurgitation of food	0.1 (0.6)	0.1 (0.6)	0.1 (0.6)	0.3 (1.0)	0.0 (0.0)
17	Denial of threat to health caused by underweight	2.5 (1.5)	3.0 (1.1)	3.2 (1.0)	0.3 (1.1)	1.9 (1.6)
	II General Psychopathology (Gen Psy)	**1.8 (0.8)**	**1.8 (0.8)**	**2.1 (0.8)**	**1.6 (0.7)**	**1.6 (0.8)**
66	Reduced self-confidence in performance	2.6 (1.2)	2.6 (1.2)	3.0 (1.2)	2.6 (1.2)	2.2 (1.3)
65	Irrational depressive thoughts and dysfunctional cognitive schemata	2.6 (1.2)	2.6 (1.3)	3.0 (1.0)	2.4 (1.3)	2.1 (1.3)
64	Depressed mood	2.7 (1.1)	2.8 (1.0)	3.0 (0.9)	2.5 (1.1)	2.2 (1.2)
67	Reduction of self-esteem	3.0 (1.1)	3.0 (1.1)	3.2 (1.1)	3.0 (0.9)	2.7 (1.2)

4	Anxieties	2.1 (1.2)	2.1 (1.2)	2.3 (1.1)	1.4 (1.0)	2.0 (1.2)
6	Sleep disturbances	2.2 (1.4)	2.1 (1.4)	2.6 (1.3)	2.1 (1.4)	1.9 (1.4)
62	Feelings of insufficiency and helplessness	2.5 (1.4)	2.4 (1.4)	2.9 (1.2)	2.5 (1.3)	2.2 (1.5)
58	Obsessive thoughts	1.5 (1 3)	1.4 (1.3)	1.8 (1.3)	0.9 (1.0)	1.6 (1.2)
68	Suicidal thoughts	1.5 (1.4)	1.3 (1.5)	1.9 (1.5)	1.1 (1.3)	1.2 (1.3)
5	Phobias	1.4 (1.2)	1.3 (1.2)	1.6 (1.3)	1.1 (1.0)	1.4 (1.2)
56	Obsessive-compulsive checking	1.4 (1.2)	1.3 (1.2)	1.6 (1.3)	0.9 (1.0)	1.5 (1.2)
69	Suicidal acts	0.5 (1.1)	0.3 (0.8)	0.9 (1.3)	0.3 (0.9)	0.4 (0.8)
70	Other autoaggressive behavior	0.7 (1.1)	0.5 (1.0)	0.9 (1.3)	0.5 (1.0)	0.6 (1.1)
57	Obsessions about cleanliness	1.0 (1.2)	0.9 (1.3)	1.3 (1.3)	0.8 (1.1)	1.0 (1.1)
	III Sexuality and Social Integration (SexSoc)	**2.1 (0.8)**	**2.2 (0.7)**	**2.4 (0.8)**	**1.7 (0.7)**	**1.7 (0.8)**
78	Sexual behavior	1.3 (1.7)	1.6 (1.7)	1.4 (1.8)	0.6 (1.4)	1.1 (1.5)
76	Sexual anxieties	1.9 (1.6)	1.9 (1.6)	2.3 (1.6)	1.4 (1.5)	1.5 (1.5)
79	Leisure activities	2.4 (1.5)	2.5 (1.4)	2.8 (1.3)	2.0 (1.5)	1.9 (1.6)
82	Social withdrawal and avoidance of contacts	2.3 (1.4)	2.3 (1.3)	2.7 (1.3)	1.9 (1.3)	1.7 (1.4)
75	Reduced sexual desire	2.4 (1.6)	2.6 (1.6)	2.8 (1.5)	1.7 (1.6)	1.8 (1.6)
77	Partner relationship	1.3 (1.3)	1.5 (1.4)	1.4 (1.3)	1.1 (1.2)	1.0 (1.0)
81	Quality of social contacts	1.3 (1.6)	1.3 (1.6)	1.6 (1.6)	1.1 (1.7)	1.1 (1.5)
83	Denial of illness	2.9 (1.2)	3.3 (1.0)	3.1 (1.1)	2.3 (1.3)	2.5 (1.1)
84	Global evaluation of symptoms	2.9 (0.9)	3.2 (0.8)	3.2 (0.8)	2.2 (0.5)	2.3 (0.8)
80	Extent of social contacts outside of family	1.8 (1.4)	1.9 (1.3)	2.3 (1.4)	1.1 (1.1)	1.4 (1.4)
53	Objective impairment at school / at work / with household	2.0 (1.4)	2.3 (1.4)	2.4 (1.4)	1.6 (1.2)	1.5 (1.2)
63	Reduced perception of internal stimuli	2.5 (1.3)	2.6 (1.3)	2.9 (1.0)	2.3 (1.4)	2.0 (1.4)
	IV Bulimic Symptoms (Bul)	**1.1 (1.1)**	**0.3 (0.4)**	**1.2 (1.1)**	**2.9 (0.6)**	**1.0 (1.0)**
21a	Binging (objectively)	0.9 (1.2)	0.2 (0.4)	1.0 (1.2)	2.7 (0.7)	0.9 (1.2)
23	Average frequency of marked eating binges (3 months)	0.8 (1.3)	0.0 (0.2)	0.8 (1.3)	3.1 (0.8)	0.6 (1.2)
24	Frequency of marked eating binges (6 months)[1]	0.7 (1.2)	0.0 (0.2)	0.7 (1.2)	2.7 (1.0)	0.6 (1.2)
21b	Binging (subjectively)	1.5 (1.5)	0.8 (1.2)	1.8 (1.6)	2.6 (1.2)	1.3 (1.4)

26	Loss of control with regard to eating	0.8 (1.3)	0.1 (0.3)	0.9 (1.4)	2.8 (0.7)	0.7 (1.2)
28	Feelings of distress regarding binge eating	0.9 (1.4)	0.1 (0.4)	1.0 (1.5)	3.1 (1.1)	0.7 (1.3)
35	Self-induced vomiting	1.2 (1.6)	0.1 (0.5)	1.5 (1.7)	3.1 (1.3)	1.1 (1.6)
34	Craving for food	1.9 (1.4)	1.3 (1.4)	1.9 (1.5)	2.8 (1.2)	2.1 (1.2)
22	Course of binge eating over time	0.7 (1.3)	0.1 (0.3)	0.8 (1.3)	2.8 (1.1)	0.6 (1.2)
	V Inappropriate Compensatory Behaviors to Counteract Weight Gain, Fasting, Substance Abuse (Compensatory behavior)	**0.9 (0.5)**	**0.6 (0.2)**	**1.1 (0.5)**	**0.9 (0.4)**	**0.9 (0.5)**
38	Appetite suppressants	0.8 (1.3)	0.1 (0.5)	1.1 (1.4)	1.1 (1.4)	0.9 (1.4)
37	Diuretics	0.3 (0.9)	0.0 (0.0)	0.6 (1.2)	0.3 (0.8)	0.4 (0.9)
41	Excessive fasting	1.1 (1.4)	0.9 (1.4)	1.5 (1.5)	0.9 (0.9)	1.0 (1.3)
72	Tranquillizers with potential for dependency	0.4 (0.8)	0.2 (0.6)	0.7 (1.0)	0.3 (0.7)	0.4 (0.8)
73	Illegal drugs	0.8 (1.0)	0.5 (0.7)	0.9 (1.1)	1.3 (1.0)	1.0 (1.0)
36	Laxative abuse	1.0 (1.4)	0.1 (0.3)	1.9 (1.5)	1.0 (1.3)	0.9 (1.4)
11	Dieting or fasting - quantitative food reduction	3.7 (0.7)	3.8 (0.5)	3.9 (0.5)	3.4 (0.9)	3.5 (0.8)
71	Alcohol use/ abuse	1.2 (1.0)	0.8 (0.8)	1.3 (1.2)	1.6 (0.9)	1.3 (0.8)
39	Medication to increase thyroid metabolism	0.1 (0.4)	0.0 (0.0)	0.1 (0.6)	0.0 (0.0)	0.0 (0.0)
43	Enemas	0.1 (0.5)	0.0 (0.0)	0.3 (0.8)	0.0 (0.0)	0.1 (0.5)
44	Ipecac	0.1 (0.4)	0.0 (0.1)	0.1 (0.6)	0.0 (0.0)	0.1 (0.2)
	VI Atypical Binges (Atyp. Binge)	**0.6 (0.8)**	**0.2 (0.4)**	**0.5 (0.8)**	**1.2 (0.8)**	**0.6 (0.9)**
31	Average number of marked atypical eating binges (3 months)	0.3 (0.8)	0.1 (0.3)	0.3 (0.8)	0.6 (0.8)	0.4 (1.0)
32	Frequency of marked atypical eating binges (6 months)[1]	0.3 (0.7)	0.1 (0.3)	0.3 (0.7)	0.6 (0.8)	0.4 (0.9)
29	Atypical eating binges	0.5 (1.0)	0.1 (0.5)	0.4 (0.9)	1.3 (1.5)	0.5 (1.0)
33	Eating more when feeling stressed or overburdend	1.2 (1.4)	0.7 (1.2)	1.3 (1.5)	2.3 (1.3)	1.3 (1.4)
30	Feeling comfortably full after binges	0.2 (0.8)	0.1 (0.5)	0.2 (0.7)	0.6 (1.3)	0.3 (0.9)
	Total Score	**1.7 (0.5)**	**1.6 (0.4)**	**2.0 (0.4)**	**1.7 (0.4)**	**1.5 (0.5)**

M=Mean; SD=Standard deviation
[1] Item is reported additonally (not included in any subscale)